arXiv:1604.00199v2 [math.QA] 29 Aug 2016

THE NODAL CUBIC IS A QUANTUM HOMOGENEOUS SPACE

ULRICH KRÄHMER AND ANGELA TABIRI

Abstract. The cusp was recently shown to admit the structure of a quantum homogeneous space, that is, its coordinate ring B can be embedded as a right coideal subalgebra into a Hopf algebra A such that A is faithfully flat as a B-module. In the present article such a Hopf algebra A is constructed for the coordinate ring B of the nodal cubic, thus further motivating the question which affine varieties are quantum homogeneous spaces.

Just as quantum groups (Hopf algebras) generalise affine algebraic groups, quantum homogeneous spaces as studied e.g. in [2, 7, 11, 14–19, 21] generalise affine varieties with a transitive action of an algebraic group:

Definition. A *quantum homogeneous space* is a right coideal subalgebra B of a Hopf algebra A which is faithfully flat as a left B-module.

There is also an analytic theory of transitive or more generally ergodic actions of compact or locally compact quantum groups, see e.g. [8, 13] and the references therein.

The best studied examples are deformation quantisations of affine homogeneous spaces, and in particular Podleś' quantum 2-spheres [20]. However, even if A is noncommutative, B can be a commutative algebra, so a natural question to ask is:

Question. *Which affine varieties are quantum homogeneous spaces?*

In the operator algebraic setting, the analogue of this question has been raised and studied for example in [12]. Here we consider the purely algebraic setting, working over an algebraically closed field k of characteristic 0.

Unlike a homogeneous space, an affine variety which is a quantum homogeneous space can be singular as the example of the cusp shows (see [15, Section 2.11] and [10, Construction 1.2]). Note that by the results from [9], noncommutative Hopf algebras coacting on commutative algebras are quite restricted. Still, we conjecture that every plane curve can be given the structure of a quantum homogeneous space, and our aim here is to point this out for the nodal cubic given by the equation $y^2 = x^2 + x^3$:

Theorem. *Fix $(q, p) \in k^2$ satisfying $p^2 = q^2 + q^3$. Then the unital associative k-algebra A with generators x, y, a, a^{-1}, b satisfying the relations*

$$aa^{-1} = a^{-1}a = 1, \qquad y^2 = x^2 + x^3, \qquad b^2 = a^3,$$

$$ba = ab, \qquad ya = ay, \qquad bx = xb, \qquad yx = xy, \qquad by = -yb + 2pb^2,$$

$$a^2x = -xa^2 - axa - a^2 + (1 + 3q)\, a^3,$$

$$ax^2 = -ax - xa - x^2a - xax + (2 + 3q)qa^3.$$

admits a Hopf algebra structure whose coproduct Δ, counit ε and antipode S satisfy

$$\Delta(x) = 1 \otimes (x - qa) + x \otimes a, \qquad \Delta(y) = 1 \otimes (y - pb) + y \otimes b,$$
$$\Delta(a) = a \otimes a, \quad \Delta(b) = b \otimes b, \quad \varepsilon(x) = q, \quad \varepsilon(y) = p, \quad \varepsilon(a) = \varepsilon(b) = 1,$$
$$S(x) = q - (x - q)\, a^{-1}, \quad S(y) = p - (y - p)\, b^{-1}, \quad S(a) = a^{-1}, \quad S(b) = b^{-1}.$$

Furthermore, the right coideal subalgebra $B \subset A$ generated by x, y is the coordinate ring of the nodal cubic, and A is free and in particular faithfully flat as a B-module.

Observe that the commutation relations in A are chosen in such a way that

$$(y - pb)^2 = y^2 - p^2 b^2, \qquad (x - qa)^2 + (x - qa)^3 = x^2 + x^3 - (q^2 + q^3)a^3$$

so that

$$(y - pb)^2 = (x - qa)^2 + (x - qa)^3.$$

Thus informally speaking each coordinate on the curve becomes perturbed by some additional group-like "quantum coordinate" and the perturbed coordinates still satisfy the defining relation of the curve. As $x - qa$ and $y - pb$ are twisted primitive, the Hopf algebra A is generated by group-likes and twisted primitives. This implies:

Proposition. *The Hopf algebra A is pointed.*

The point (q, p) on the curve is the point the quantum orbit of which it is presented as. To use these observations as starting point of a general study of quantum homogeneous space structures on affine varieties seems a promising future research direction.

The proof of the theorem consists of a straightforward (albeit tedious) verification that the formulas for the coproduct, counit and antipode are compatible with the defining relations of A, followed by a similarly straigthforward application of Bergman's diamond lemma [1] yielding a vector space basis of A that implies the freeness over B:

Proposition. *The set*

$$\{x^i y^j (ax)^l a^m b^n \mid i, l \in \mathbb{N}, j \in \{0, 1\}, m \in \mathbb{Z}, n \in \{0, 1\}\}$$

is a vector space basis of A, and the GK-dimension of A equals 3.

Using this basis, one also easily observes that like the nonstandard Podleś spheres, the algebra extension $B \subset A$ is an example of a coalgebra Galois extension [3–6] rather than of a Hopf-Galois extension: the coalgebra is $C := A/B^+A$, where $B^+ := B \cap \ker \varepsilon$. The canonical projection $\pi : A \to C$ defines a left C-coaction

$$\lambda \colon A \to C \otimes A, \quad f \mapsto f_{(-1)} \otimes f_{(0)} := \pi(f_{(1)}) \otimes f_{(2)}$$

and we have (as a consequence of the faithful flatness of A over B)

$$B = \{f \in A \mid f_{(-1)} \otimes f_{(0)} = \pi(1) \otimes f\}.$$

That C is not a Hopf algebra quotient of A follows from $B^+A \neq AB^+$ (cf. [18, Lemma 1.4]); for example, we have

$$AB^+ \ni a^2(x - q) = -xa^2 - axa - (1 + q)a^2 + (1 + 3q)a^3 \notin B^+A.$$

We finally remark that some properties of the algebra A are better understood when using a slightly different set of generators: if we abbreviate

$$c := 3x - (1+3q)a + 1, \quad d := 3y - 6pb, \quad e := ac + rca$$

where r is a primitive 6th root of 1 (so that $r + r^{-1} = 1$), then the defining relations of A in terms of the generators $a^{\pm 1}, b, c, d, e$ read

$$aa^{-1} = a^{-1}a = 1, \quad ab = ba, \quad ac + rca = e, \quad ad = da, \quad ae + r^{-1}ea = 0,$$
$$bc = cb, \quad bd = -db, \quad be = eb, \quad b^2 = a^3, \quad cd = dc, \quad r^{-1}ce + ec = 3(a - a^3),$$
$$de = ed, \quad 3d^2 = c^3 - 3c + 2 + (1+3q)(-2 + 6q + 9q^2)a^3.$$

Using these generators, one easily verifies for example:

Proposition. *The units in A are of the form $\alpha a^i b^j$, $\alpha \in k$, $i \in \mathbb{Z}$, $j \in \{0, 1\}$.*

Acknowledgements. We thank Ken Brown and Tomasz Brzeziński for helpful comments on a first draft of this paper. A.T. is funded by a Faculty for the Future Fellowship of the Schlumberger Foundation.

References

[1] George M. Bergman, *The diamond lemma for ring theory*, Adv. in Math. **29** (1978), no. 2, 178–218, DOI 10.1016/0001-8708(78)90010-5. MR506890

[2] Ken Brown and Paul Gilmartin, *Quantum homogeneous spaces of connected Hopf algebras*, J. Algebra **454** (2016), 400–432, DOI 10.1016/j.jalgebra.2016.01.030. MR3473433

[3] Tomasz Brzeziński, *On modules associated to coalgebra Galois extensions*, J. Algebra **215** (1999), no. 1, 290–317, DOI 10.1006/jabr.1998.7738. MR1684158

[4] Tomasz Brzeziński and Piotr M. Hajac, *Coalgebra extensions and algebra coextensions of Galois type*, Comm. Algebra **27** (1999), no. 3, 1347–1367, DOI 10.1080/00927879908826498. MR1669095

[5] ———, *The Chern-Galois character*, C. R. Math. Acad. Sci. Paris **338** (2004), no. 2, 113–116, DOI 10.1016/j.crma.2003.11.009 (English, with English and French summaries). MR2038278

[6] Tomasz Brzeziński and Shahn Majid, *Coalgebra bundles*, Comm. Math. Phys. **191** (1998), no. 2, 467–492, DOI 10.1007/s002200050274. MR1604340

[7] Kenny De Commer, *On the construction of quantum homogeneous spaces from *-Galois objects*, Algebr. Represent. Theory **15** (2012), no. 4, 795–815, DOI 10.1007/s10468-011-9265-7. MR2944442

[8] Kenny De Commer and Makoto Yamashita, *Tannaka-Kreĭn duality for compact quantum homogeneous spaces. I. General theory*, Theory Appl. Categ. **28** (2013), No. 31, 1099–1138. MR3121622

[9] Pavel Etingof, Debashish Goswami, Amab Mandal, and Chelsea Walton, *Hopf coactions on commutative algebras generated by a quadratically independent comodule*, Comm. Algebra (to appear).

[10] Kenneth R. Goodearl and James J. Zhang, *Noetherian Hopf algebra domains of Gelfand-Kirillov dimension two*, J. Algebra **324** (2010), no. 11, 3131–3168, DOI 10.1016/j.jalgebra.2009.11.001. MR2732991

[11] István Heckenberger and Stefan Kolb, *Differential calculus on quantum homogeneous spaces*, Lett. Math. Phys. **63** (2003), no. 3, 255–264, DOI 10.1023/A:1024496304742. MR1992890

[12] Huichi Huang, *Faithful compact quantum group actions on connected compact metrizable spaces*, J. Geom. Phys. **70** (2013), 232–236, DOI 10.1016/j.geomphys.2013.03.027. MR3054297

[13] Paweł Kasprzak and Piotr M. Sołtan, *Embeddable quantum homogeneous spaces*, J. Math. Anal. Appl. **411** (2014), no. 2, 574–591, DOI 10.1016/j.jmaa.2013.07.084. MR3128415

[14] Stefan Kolb and Jacopo Pellegrini, *Braid group actions on coideal subalgebras of quantized enveloping algebras*, J. Algebra **336** (2011), 395–416, DOI 10.1016/j.jalgebra.2011.04.001. MR2802552

[15] Ulrich Krähmer, *On the Hochschild (co)homology of quantum homogeneous spaces*, Israel J. Math. **189** (2012), 237–266, DOI 10.1007/s11856-011-0168-4. MR2931396

[16] Liyu Liu and Quanshui Wu, *Twisted Calabi-Yau property of right coideal subalgebras of quantized enveloping algebras*, J. Algebra **399** (2014), 1073–1085, DOI 10.1016/j.jalgebra.2013.10.019. MR3144627

[17] ———, *Rigid dualizing complexes over quantum homogeneous spaces*, J. Algebra **353** (2012), 121–141, DOI 10.1016/j.jalgebra.2011.12.007. MR2872439

[18] Ernst F. Müller and Hans.-Jürgen Schneider, *Quantum homogeneous spaces with faithfully flat module structures*, Israel J. Math. **111** (1999), 157–190, DOI 10.1007/BF02810683. MR1710737

[19] Réamonn Ó Buachalla, *Noncommutative complex structures on quantum homogeneous spaces*, J. Geom. Phys. **99** (2016), 154–173, DOI 10.1016/j.geomphys.2015.10.003. MR3428362

[20] Piotr Podleś, *Quantum spheres*, Lett. Math. Phys. **14** (1987), no. 3, 193–202, DOI 10.1007/BF00416848. MR919322

[21] Hans-Jürgen Schneider, *Principal homogeneous spaces for arbitrary Hopf algebras*, Israel J. Math. **72** (1990), no. 1-2, 167–195, DOI 10.1007/BF02764619. Hopf algebras. MR1098988

University of Glasgow, School of Mathematics and Statistics, 15 University Gardens, G12 8QW Glasgow, UK

E-mail address: ulrich.kraehmer@glasgow.ac.uk,a.tabiri1@research.gla.ac.uk